



POSTER PRESENTATION

Open Access

Anti-IL1 β -monoclonal antibody in a patient with Muckle-Wells Syndrome and renal transplantation - five years experience

B Kortus-Götze*, J Hoyer

From 8th International Congress of Familial Mediterranean Fever and Systemic Autoinflammatory Diseases Dresden, Germany. 30 September - 3 October 2015

Background

The Muckle-Wells syndrome (MWS) is a rare inherited disease and belongs to the group of cryopyrin-associated periodic syndromes (CAPS). Recurrent fever attacks, myalgia, arthralgia, urticarial rash, headache, conjunctivitis, sensorineural deafness and a severe fatigue syndrome are the typical symptoms of MWS. Due to an unregulated production of IL1 a continuous formation of serum amyloid leads ultimately to the development of AA-amyloidosis, which is life-threatening and in some cases the fatal complication of MWS.

Objectives

The anti-IL1 β -monoclonal antibody canakinumab has been introduced as a specific therapy in patient with MWS with normal and impaired renal function. There are no long term data about the effect from canakinumab in patients with renal transplantation.

Methods

Here we report the five years follow up on a 36-year old female patient with Muckle-Wells syndrome and biopsy proven systemic AA amyloidosis and end stage renal disease. After renal transplantation therapy with canakinumab subcutaneously in a dosage of 150 mg every eight weeks was continued in combination with the immunosuppressive therapy.

Results

Before and after renal transplantation the patient had a very good response to canakinumab with low activity in inflammation markers with an improved quality of life. Over the period of five years the triple immunosuppressive

therapy (CSA, MMF, Prednisone) in combination with canakinumab has had no negative effect on activity of MWS and no pharmacological interactions between medications were observed. Even five years after renal transplantation, the patient remains an excellent kidney function without proteinuria. There are no signs of recurrence of AA-amyloidosis in the transplanted kidney.

Conclusions

According to our data, treatment with different immunomodulators in patients with Muckle-Wells syndrome and renal transplantation is safe, feasible and without severe side effects also over a longer time.

Published: 28 September 2015

doi:10.1186/1546-0096-13-S1-P147

Cite this article as: Kortus-Götze and Hoyer: Anti-IL1 β -monoclonal antibody in a patient with Muckle-Wells Syndrome and renal transplantation - five years experience. *Pediatric Rheumatology* 2015 13(Suppl 1):P147.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



University of Marburg, Nephrology, Marburg, Germany



© 2015 Kortus-Götze and Hoyer This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.