



ORAL PRESENTATION

Open Access

Overexpression of CREM α accelerates onset and severity of auto-immune mediated disease in a murine model of lupus

Kim Ohl^{1*}, Ralph Lippe², Eva Verjans¹, Nora Honke¹, Karin Maschke-Neuss¹, Norbert Wagner¹, Johannes Roth², Klaus Tenbrock¹

From 18th Pediatric Rheumatology European Society (PReS) Congress
Bruges, Belgium. 14-18 September 2011

The transcription factor cAMP response element modulator (CREM) is a widely expressed transcriptional repressor which is important for the termination of the T cell immune response. CREM α is overexpressed in SLE (Systemic lupus erythematosus) T cells and is supposed to be a key player in orchestrating the transcriptional program of SLE T cells by targeting T cell-relevant genes. To explore the relevance of CREM α in vivo we used a well-established murine lupus model, which is characterized by the introduction of a mutation in the CD95 (Fas) locus. We generated a transgenic mouse with a selective overexpression of CREM α in T cells and introduced a Fas $-/-$ phenotype into the CREM α transgenic mice. CREM α transgenic Fas $-/-$ mice developed a severe lymphadenopathy and splenomegaly as early as 8 weeks of age, while the wildtype Fas $-/-$ mice did not at this early age. Lymphadenopathy and splenomegaly is paralleled by a massive expansion of pathogenic CD3+CD4-CD8- double negative T cells. Furthermore T cells of CREM α transgenic Fas $-/-$ mice show an enhanced production of IL-21 and IL-17, which are hallmark cytokines of highly inflammatory Th17 cells. Vice versa percentages of regulatory T cells are reduced. The enhanced occurrence of aberrant and inflammatory T cells further leads to increased B cell activation, increased anti-DNA antibody titers and finally shortened life expectation of these mice.

Our experiments are the proof of principle for a critical amplifying role of CREM α in autoimmune prone conditions like SLE.

Author details

¹Klinik für Kinder und Jugendmedizin, Klinikum der RWTH Aachen, Germany.

²Institut für Immunologie, Universität Münster, Germany.

Published: 14 September 2011

doi:10.1186/1546-0096-9-S1-O3

Cite this article as: Ohl et al.: Overexpression of CREM α accelerates onset and severity of auto-immune mediated disease in a murine model of lupus. *Pediatric Rheumatology* 2011 **9**(Suppl 1):O3.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: kohl@ukaachen.de

¹Klinik für Kinder und Jugendmedizin, Klinikum der RWTH Aachen, Germany
Full list of author information is available at the end of the article