



POSTER PRESENTATION

Open Access

A validation of diagnostic score for molecular analysis of hereditary autoinflammatory syndromes with periodic fever in Turkish children

Erkan Demirkaya^{1*}, Cengizhan Acikel², Isabella Ceccherini³, Seza Özen⁴, Abdulkaki Karaoglu¹, Ismail Dursun⁵, Betül Sozeri⁶, Yelda Bilginer⁴, Zubeyde Gunduz⁷, Yusuf Tunca⁸, Katerina Oikonomaki³, Ioanna Varela³, Francesco Caroli³, Maria Pia Sormani⁹, Marco Gattorno¹⁰

From 18th Pediatric Rheumatology European Society (PReS) Congress Bruges, Belgium. 14-18 September 2011

Objective

In the present study, we validated a set of variables that predict the presence of a mutation in one of the 3 genes known to be associated with the common monogenic autoinflammatory diseases.

Methods

A total of 93 consecutive patients with a clinical history of periodic fever were screened for mutations in the *MVK*, *TNFRSF1A*, and *MEFV* genes, and detailed clinical information was collected prospectively. For autosomal-recessive diseases (MKD and FMF), only homozygous or compound heterozygous patients were accepted as genetically positive. The developed diagnostic score was validated in an independent set of 93 patients.

Results

93 patients whose suspected diagnosis was one of the autoinflammatory diseases were enrolled. The mean age at onset was 58,64+40,69 months (54 F, 39 M;). A total of 25 patients with periodic or recurrent fevers did not display any mutation in the 3 genes examined. Of these 68 patients with positive findings on genetic testing, 30 of them had a mutation only in one allele and 38 had mutations in two alleles for *MEFV* gene. In addition, we identified 4 low-penetrance R92Q mutations for *TRAPS* and 3 patients, who were compound heterozygotes for the *MVK* gene. The diagnostic score revealed moderate sensitivity (68%) and specificity (58%) for discriminating

patients who were genetically positive from those who were negative.

Conclusion

The suggested set of variables may be used to decide on genetic screening for patients with periodic fever syndromes. However, further studies are needed to improve the sensitivity of the suggested score.

Author details

¹Gulhane Military Medical Academy, Pediatric Nephrology and Rheumatology Unit, Ankara, Turkey. ²Gulhane Military Medical Faculty, Department of Public Health, Division of Epidemiology, Ankara, Turkey. ³Laboratorio di Genetica Molecolare, Istituto Giannina Gaslini, Genova, Italy. ⁴Hacettepe University Faculty of Medicine, Department of Pediatrics, Pediatric Nephrology and Rheumatology Unit, Ankara, Turkey. ⁵Kayseri Education and Research Hospital, Kayseri, Turkey. ⁶Ege University Faculty of Medicine Department of Pediatrics, Division of Nephrology, Izmir, Turkey. ⁷Erciyes University Medical School, Pediatric Nephrology & Rheumatology Unit, Kayseri, Turkey. ⁸Gulhane Military Medical Academy, Department of Medical Genetics, Ankara, Turkey. ⁹University of Genoa, Genoa, Italy. ¹⁰Giannina Gaslini Institute, 2nd Division of Paediatrics, Genoa, Italy.

Published: 14 September 2011

doi:10.1186/1546-0096-9-S1-P298

Cite this article as: Demirkaya et al.: A validation of diagnostic score for molecular analysis of hereditary autoinflammatory syndromes with periodic fever in Turkish children. *Pediatric Rheumatology* 2011 **9**(Suppl 1):P298.

* Correspondence: dottore_erkana@yahoo.com

¹Gulhane Military Medical Academy, Pediatric Nephrology and Rheumatology Unit, Ankara, Turkey

Full list of author information is available at the end of the article